

Giving Diastat to a Child:

www.diastat.com

Before giving Diastat:


1. Read instructions and understand how to give the Diastat

2. Reviewed when to give it with the doctor, if a dose should be repeated, and when to call for emergency help.


Please do not give DIASTAT® AcuDial™ until:

1. You have confirmed:
 - Prescribed dose is visible and if known, is correct
 - Green “ready” band is visible
2. Know how to dispose of the medicine.


CHILD ADMINISTRATION INSTRUCTIONS




1 Put person on their side where they can't fall.




4 Push up with thumb and pull to remove cap from syringe. **Be sure seal pin is removed with the cap.**




5 Lubricate rectal tip with lubricating jelly.




8 Separate buttocks to expose rectum.




9 Gently insert syringe tip into rectum. **Note: rim should be snug against rectal opening.**




12 Slowly count to 3 while holding buttocks together to prevent leakage.




11 Slowly count to 3 before removing syringe from rectum.




10 Slowly count to 3 while gently pushing plunger in until it stops.



13 **ONCE DIASTAT™ IS GIVEN** Keep person on the side facing you, note any change in breathing, pulse, and continue to observe.



3 Get syringe. **Note: seal pin is attached to the cap.**



6 Turn person on side facing you.

2 Get medicine.

7 Bend upper leg forward to expose rectum.

10...11...12...13 **COUNT OUT LOUD TO THREE...**

CALL FOR HELP IF ANY OF THE FOLLOWING OCCUR

- Seizure(s) continues 15 minutes after giving DIASTAT™ or per the doctor's instructions
- Seizure behavior is different from other episodes
- You are alarmed by the frequency or severity of the seizure(s)
- You are alarmed by the color or breathing of the person
- The person is having unusual or serious problems

Local emergency number: _____ Doctor's number: _____
(Please be sure to note if your area has 911)

Information for emergency squad: Time DIASTAT™ given: _____ Dose: _____

DIASTAT™ Indication
DIASTAT™ AcuDial™ (diazepam rectal gel) is a gel formulation of diazepam intended for rectal administration in the management of seizure. Intended for use with patients with a history of Lennox-Gastaut syndrome (LGS) and patients with a history of diazepam to control bouts of increased seizure activity, for patients 2 years and older.

Important Safety Information
In clinical trials with DIASTAT™, the most frequent adverse event was somnolence (23%). Less frequent adverse events reported were dizziness, headache, pain, vasodilation, diarrhea, ataxia, euphoria, incoordination, asthma, rash, abdominal pain, nervousness, and thrush (1%-5%).

D955-0308

Diastat®
(diazepam rectal gel)

Diastat AcuDial™
(diazepam rectal gel)

DISPOSAL INSTRUCTIONS ON REVERSE SIDE **62**